

# Abstract of Priority Documents

WPIDS COPYRIGHT 2003 THOMSON DERWENT on STN

AN 2001-356049 [37] WPIDS

CR 2001-356050 [37]

DNN N2001-258683 DNC C2001-110489

TI Detecting multiple groups such as kinases or kinase substrates in test sample, comprises separating proteins in sample to produce one-dimensional array, contacting array with antibodies and detecting antibodies bound to groups.

DC B04 D16 S03

IN PELECH, S

PA (UYBR-N) UNIV BRITISH COLUMBIA

CYC 25

PI WO 2001038877 A2 20010531 (200137)\* EN 61p G01N033-573

RW: AT BE CH CY DE DK ES FI FR GB GR IE IT LU MC NL PT SE TR

W: AU CA JP NZ US

CA 2290335 A1 20010519 (200141) EN C12Q001-48

CA 2290204 A1 20010522 (200143) EN C12Q001-48

AU 2001016843 A 20010604 (200153) G01N033-573

EP 1234184 A2 20020828 (200264) EN G01N033-573

R: AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE TR

ADT WO 2001038877 A2 WO 2000-CA1377 20001117; CA 2290335 A1 CA 1999-2290335

19991119; CA 2290204 A1 CA 1999-2290204 19991122; AU 2001016843 A AU

2001-16843 20001117; EP 1234184 A2 EP 2000-979296 20001117, WO 2000-CA1377

20001117

FDT AU 2001016843 A Based on WO 2001038877; EP 1234184 A2 Based on WO

2001038877

PRAI US 2000-216357P 20000705; CA 1999-2290335 19991119; CA 1999-2290204

19991122

IC ICM C12Q001-48; G01N033-573

ICS C12Q001-42; G01N033-535

AB WO 200138877 A UPAB: 20021031

NOVELTY - Detecting (M) multiple groups such as kinases or kinase substrates in a test sample comprising:

(1) electrophoretically separating proteins in the sample to produce a one-dimensional array of proteins so separated;

(2) contacting the array with antibodies such as anti-kinase (substrate) antibodies; and

(3) detecting the presence of antibodies bound to kinase (substrate) groups in the array, is new.

USE - (M) is useful for detecting multiple groups such as kinases or kinase substrates in a test sample (claimed). (M) is useful for the discovery of new protein kinases, to detect unknown proteins which can cross-react with kinase-specific antibodies, and to identify known proteins and detect new proteins that may bind with high affinity to a target protein.

ADVANTAGE - (M) has many advantages over standard 2-dimensional (2D) gel proteomic methods. (M) can be applied to any cell or tissue samples, and no prelabeling with radioisotopes is necessary, because kinase detection is based on immunoreactivity. (M) can be adapted for wide scale diagnostic applications because patterns of protein kinase expression are stable for periods of up to six hours before an organ is subjected to fractionation and freezing, providing the organ is stored during this time over ice. This procedure may be carried out within two days from start to finish when compared to the 2D gel electrophoresis which is extremely laborious, much more difficult to render and takes at least twice a time. (M) provides ability to compare multiple samples side by side, whereas the 2D gel can be used for a single sample. An advantage of (M) over other methods to examine protein-protein interactions such as the yeast two-hybrid method, is that it can detect interactions that are affected by the state of post-translation regulation of these proteins, such as their phosphorylation state.

Dwg. 0/6

FS CPI EPI

FA AB; DCN